

Adapting Tracking Techniques Used on Least Terns to Coastal Species of Concern

Jennifer H. Stucker
Mark H. Sherfy

Northern Prairie Wildlife
Research Center
Jamestown, ND USA

U.S. Department of the Interior
U.S. Geological Survey




Background Telemetry

- Antenna

- Receiver

- Transmitter

- Recent developments change traditional “receiver” and “transmitter”




Background "Transmitters"

	Broadcast	Size	Benefits	Drawbacks
Radio-VHF <i>Signal</i>	1 freq. to 1 transmitter 60 freq. \Rightarrow 60 transmitters	≥ 0.5 g	<ul style="list-style-type: none"> • Simple & efficient • extras • Small 	<ul style="list-style-type: none"> • simple, • need receiving method • Z=extra weight
<i>Coded</i>	1 freq. \Leftrightarrow 10X transmitters	≥ 1 g	<ul style="list-style-type: none"> • Simple & efficient • Simultaneous detection of individuals 	<ul style="list-style-type: none"> • simple, • need receiving method <i>and limited</i> manual relocation • Z=extra weight
Satellite (PTT)	To satellite	≥ 17 g	<ul style="list-style-type: none"> • Real time; • can obtain Z • no receivers 	<ul style="list-style-type: none"> • Heavier, • power hungry, • depend on satellites • Z \pm err (10-50 m)
GPS	none – stored in unit; vhf	≥ 22 g	<ul style="list-style-type: none"> • Real time; • can obtain Z • no receivers 	<ul style="list-style-type: none"> • Have to retrieve to get data – or batch download via satellite. • depend on satellites • Z \pm err (10-50 m)

Background Receiving - VHF

- **Fixed or mobile Bi- & Tri+ angulation**

Requires "stationary" critter



- **Aircraft flight**

Altitude + visibility restriction
Infrequent relocation

- **Continuous stationary logging**


Presence-absence limited area
Frequent relocation
Omni-directional
Multi-directional

Background Choices?


- Chose right equipment by question.
- “Transmitter” size limited by size of animal.
 - For birds $<3\%$ weight
- Possible to make up “limitations” of VHF transmitter with receiving and monitoring choices to address research question.

Missouri River Flows

- Historical (pre-dam):
 - Pronounced spring peak in hydrograph
 - Wide, braided channel with numerous sandbars
- Modern (post-1944):
 - 6 dams (South Dakota to Montana)
 - Reservoirs
 - Flow-managed rivers
 - Navigation
 - Hydropower
 - Flood Control
 - Recreation
 - Wildlife & Fisheries



Overview Missouri River



~2,300 miles MT to MO

~700 miles channelized
Sioux City, IA to
Mississippi River

Federally Listed

- Pallid Sturgeon
- Piping Plover
- Least Tern

*"This bird is very noysey
when flying which it dose
exttreemly swift...."*

Meriwether Lewis, 1804

Overview Interior Least Tern

Sternula antillarum

Laridae

- Smallest NA tern, 45 g
- Interior population, Endangered
- Ground nesting
- 2-3 eggs
- Eats small fish
- Greatest threat – nesting habitat loss



Least Tern population status

- California population – Endangered
- Gulf and Atlantic - no Federal Listing

Overview Least Terns and Missouri River

- **Biological Opinion**

1. Emergent sandbar habitat
2. Reproductive success

- **Sandbars created for nesting habitat**

# sandbar projects	Completion year
1	2004
2	2005
1	2007-2008
3-4	2008



Overview – Least Terns on Gavins Point Reach

- **Least Tern use of natural(3) and created (3) sandbars**
- **Are there differences between created and natural sandbars?**


Multiple methods integrated to evaluate response

- Productivity & nesting habitat
- Foraging habitat
 - Fish sampling
- Behavior
- Movements



Methods

- Preliminary results presented are confined to birds nesting at RM 770 in 2006 and 2007, and their use of the river.



Methods

Behavior

- Colony scan samples
⇒ activity budgets
- Flying and foraging on river
- Spot map locations

Movements

- Telemetry
- Assessing Duration In/Out colony
- Evaluate frequency of use of river

Fish Sampling



Methods Radio Transmitters

- Failure with Least Terns
(Massey et al. 1988)
- Recent use other terns
- Attachment trial



Field

- Trapped on nest
- Marked
 - Bands (metal + color), transmitter, plumage color marks



Methods Telemetry

Null-peak set-up

- bi- or triangulation methods – NO

Boats surveys

- 2 – 4 surveys per week - YES



Methods Telemetry receiving

- Stationary data loggers



	<u>2006</u>	<u>Antenna</u>	<u>2007</u>	<u>Antenna</u>
colony	6	1	7	1
river	1 (5)	1	8	2-4



Summary preliminary

Based on data collected for Least Terns at RM 770



- Changes in colony behavior through season
- Differences foraging behavior and fish capture
- Variation in river use by foraging Least Terns
- Changes in amount of river used by breeding stage
- Document differences in nocturnal vs. diurnal activity via telemetry

Will these patterns hold for all natural and created sandbars?

- 2008 final field season; complete evaluation for all (~10) sandbars expected in 2009.

Would methods work in coastal waters?

Benefits


duration

frequency

timing

Challenges

- One dimension, multi-dimensional ... ?



- Land & fresh water *to* saltwater; tides
- Remote access capabilities

Acknowledgements

Funding:

US Army Corps of Engineers, Omaha District T&E section

Sally Valdes and MMS for invitation to participate

Field Assistance:

Crew Chief, boat trainer, & bird trapper: Colin Dovichin

Crew Leaders:

2006: Brandi Skone (behavior), John Campbell (telemetry), Michael Sertle (nests)

2007: Brandi Skone (behavior), John Campbell (telemetry), Mitch Johnson (fish),
Breanna Riedel (nests)

Technicians:

Alicia Andes, Amanda Bryson, Craig Davis, Michael Duchscher, Erin Ellmaker, Danielle Haak, Bryant Huso, Kaylan Kemink, Kurtis Lammers, Heidi Murray, Jessica Orr, Ashley Persinger, Chris Petersen, Dolph Prater, Laell Schulte, Jordan Wein, Brenda Jarski-Weber, Nate Emery, Janelle Feine, Mary Chris Harrison, Zach Kaiser, Jason Kottsick, Erik Schmidt, Samantha Smith, Jenny Yoo

Contact Information

Jennifer H. Stucker

USGS - Northern Prairie Wildlife Research Center

Jamestown, ND 58401

701-253-5539

jstucker@usgs.gov